

Diagnostic Challenge

On the tip of the tongue . . .

Rachel Orchard, MD*; Sheena Belisle, MD†; Rodrick Lim, MD†‡

Keywords: pediatric, rash, tongue, vesicle

CASE HISTORY

A 14-year-old male presented to the pediatric emergency department (ED) with a chief complaint of changes to his tongue. He described a 3-day history of a gradually worsening sore, swollen tongue associated with a white plaque. This was accompanied by a 3-day history of a gradually worsening left-sided facial rash that had an intermittent mild tingling sensation. He also had a 1-week history of a productive cough with yellow mucus and generalized malaise. He had been seen at a walk-in clinic 2 days prior to presentation and was prescribed amoxicillin for presumed pneumonia, which he began the same day. He denied any history of fevers, facial weakness, neck stiffness, or eye symptoms. He was an otherwise well child, with up-to-date immunizations and a past medical history of chickenpox and recurrent furuncles as a younger child.

On examination, he appeared well with the following vital signs: blood pressure 122/64 mm Hg, heart rate 73 beats per minute, respiratory rate 18 breaths per minute, temperature 36.8°C, and oxygen saturation of 99% on room air. Examination of his tongue revealed a symmetric white plaque along with ulcerative lesions on the left tongue and buccal mucosa (Figure 1). Head and neck examination revealed a left-sided facial vesicular rash, extending from the corner of the mouth to the ear (Figure 2). There were no lesions surrounding the eye or within the external auditory canal. Respiratory examination revealed a mild

right-sided wheeze. Cardiovascular, abdominal, and neurological (including cranial nerve) examinations were unremarkable.

What is the most likely diagnosis?

- a) Drug eruption
- b) Varicella zoster virus (VZV)
- c) Oral candidiasis
- d) Epstein-Barr virus
- e) Oral lichen planus




Figure 1. Leukoplakia and ulcerative lesions on tongue.

From the *Department of Family Medicine, †Department of Paediatrics, and ‡Department of Medicine, Schulich School of Medicine & Dentistry, University of Western Ontario, Children’s Health Research Institute, Children’s Hospital London Health Sciences Centre, London, ON.

Correspondence to: Dr. Rodrick Lim, Children’s Hospital London Health Sciences Centre, 800 Commissioners Road East, London, ON N6C 2V5; Email: Rod.Lim@lhsc.on.ca



Figure 2. Facial rash extending from corner of mouth to ear.

Answer: This patient's history and physical examination are consistent with a VZV infection, manifesting as herpes zoster (HZ). HZ (shingles) is caused by reactivation of latent VZV in the cranial nerves or dorsal root ganglia, and most commonly presents as an erythematous vesicular rash grouped along a sensory dermatome.¹ VZV can also manifest as a primary infection, commonly referred to as *varicella* (chickenpox), which classically presents as a diffuse vesicular rash over the face, trunk, and extremities.

HZ most commonly affects older adult patients or those with some form of immunodeficiency, but is also seen in the otherwise healthy pediatric population. A recent U.S. study² found an incidence of 112 per 100,000 person-years in children ages 0 to 17 years. HZ can be caused by both wild-type VZV and vaccine-strain VZV; however, vaccinated children have a lower incidence rate of zoster than unvaccinated children.²⁻⁴ There also appears to be a higher risk of developing zoster if varicella is contracted in the first year of life.^{3,4} Unlike the adult population, pediatric zoster is associated with minimal pain on initial presentation and is rarely complicated by postherpetic neuralgia. It most commonly affects the thoracic dermatomes, with 5% of cases involving cranial nerves.⁴

Table 1. Differential diagnosis of oral leukoplakia, and oral lesions that can mimic leukoplakia⁵⁻⁹

Neoplastic
• Squamous cell carcinoma
Infectious
• Oropharyngeal candidiasis
• Human immunodeficiency virus (HIV)
• Cytomegalovirus (CMV)
• Epstein-Barr virus (EBV)
• Herpes simplex virus (HSV)
• Varicella zoster virus (VZV)
• Coxsackie virus
Autoimmune/inflammatory
• Systemic lupus erythematosus (SLE)
• Bullous pemphigoid
• Pemphigus vulgaris
• Behcet syndrome
• Crohn disease
Miscellaneous
• Aphthous stomatitis
• Erythema multiforme +/- Stevens-Johnson syndrome
• Oral lichen planus
• Geographic tongue
• Hairy tongue

The facial rash in this case presentation was consistent with HZ, given the grouped vesicles spread along the maxillary (CN V₂) and mandibular (CN V₃) divisions of the trigeminal nerve (CN V). However, the leukoplakia remained more of a diagnostic challenge. Although the left-sided facial rash pointed us towards a diagnosis of zoster, the symmetry of the leukoplakia was puzzling; this prompted consideration of the differential diagnosis of leukoplakia (Table 1) and a review of the cranial nerve innervation of the tongue. As shown in Figure 3, the lingual nerve, a branch of the CN V₃, supplies sensory innervation to the anterior two-thirds of the tongue. This explains the potential for spread of lesions to the tongue in a VZV infection of the CN V. Anterior tongue involvement can also occur in VZV affecting the facial nerve (CN VII) and has been previously documented in case reports¹⁰⁻¹³ of the Ramsay Hunt syndrome (further discussed after the following paragraph).

In this case, on careful inspection, the ulcers were more prominent in the anterior left tongue, with secondary leukoplakia developing. A review of the literature did not reveal any published statistics on the frequency of associated leukoplakia in VZV infections of the trigeminal nerve (specifically, CN V₃). Part of this may be due to




Figure 3. Cranial nerve innervation of the tongue.

varying definitions of leukoplakia amongst clinicians (as a descriptive term versus a specific diagnosis); however, it is known that mandibular nerve involvement of VZV can result in ulcerative lesions appearing on the tongue, buccal mucosa, and gingival.¹⁴ Confluent ulcerations or vesicles can often appear similar to leukoplakia, and, in fact, there is a case report¹⁵ of VZV affecting the trigeminal nerve in which the tongue was unilaterally affected by white ulcerations, in a manner somewhat similar to this patient's presentation.

For any patient with a suspected zoster infection involving the head and neck, it is important to rule out HZ ophthalmicus (infection of the ophthalmic nerve branch, CN V₁). Evaluation for visual symptoms, lesions around the eye, and lesions around the nose (Hutchinson sign) is crucial. It is also important to look for conjunctival injection and edema, because conjunctivitis is one of the most common presentations of HZ ophthalmicus.¹⁶ If ophthalmic nerve involvement is suspected, a thorough eye exam with ophthalmology consultation is warranted.¹⁶ It is also important to evaluate the patient for facial paralysis, lesions within the

ear canal and adjacent skin, otalgia, and hearing loss. These symptoms can indicate HZ oticus (also known as *Ramsay Hunt syndrome*), which arises from VZV infection of the CN VII. The geniculate ganglion of CN VII has two major sensory branches; the first is responsible for taste sensation to the anterior tongue via the chorda tympani, and the second supplies sensation to the external auditory canal and pinna.¹⁷ Ramsay Hunt syndrome is managed with a course of acyclovir, as well as corticosteroids (such as prednisone) in severe cases.⁴

Our patient was treated as a HZ infection involving CN V₂ and V₃, and was given a 7-day course of acyclovir 800 mg five times daily. His oral antibiotics were discontinued because his chest radiograph was clear. Given his wheeze, a component of asthma was suspected and Ventolin was trialed with success. Initiation of oral antibiotics 2 days prior to presentation to the ED was likely appropriate given his productive cough, auscultatory finding presumably consistent with pneumonia, and absence of significant facial findings at that time. He was seen in follow-up by our pediatric infectious disease specialist several days after initial presentation, by which time the facial lesions were already crusting over. Swabs taken of a vesicular lesion on his face returned positive for VZV via polymerase chain reaction (PCR) genotyping, confirming our diagnosis. He was seen again in follow-up 2 weeks later, by which time the tongue lesions had resolved as well.

Competing interests: None declared.

REFERENCES

- Steiner I, Kennedy PGE, Pachner AR. The neurotropic herpes viruses: herpes simplex and varicella-zoster. *Lancet Neurol* 2007;6(11):1015-28.
- Weinmann S, Chun C, Schmid DS, et al. Incidence and clinical characteristics of herpes zoster among children in the varicella vaccine era, 2005-2009. *J Infect Dis* 2013; 208(11):1859-68.
- Wen SY, Wen-Liang L. Epidemiology of pediatric herpes zoster after varicella infection: a population-based study. *Pediatrics* 2015;135(3):e555-71.
- Feder HM, Hoss DM. Herpes zoster in otherwise healthy children. *Pediatr Infect Dis J* 2004;23(5):451-60.
- Gonsalves WC, Chi AC, Neville BW. Common oral lesions: part I. Superficial mucosal lesions. *Am Fam Phys* 2007; 75(4):501-6.
- Gonsalves WC, Chi AC, Neville BW. Common oral lesions: part II. Masses and neoplasia. *Am Fam Phys* 2007;75(4): 507-12.

7. Chi AC, Neville BW, Krayner JW, et al. Oral manifestations of systemic disease. *Am Fam Phys* 2010;82(11):1381-8.
8. Parent D. Oral viral infections. *Presse Med* 2016;45(2):195-214.
9. Siu A, Landon K, Ramos DM. Differential diagnosis and management of oral ulcers. *Semin Cutan Med Surg* 2015;34(4):171-7.
10. Lee CK, Baek BJ. Images in clinical medicine: lingual zoster. *N Engl J Med* 2011;365(18):1726.
11. Zhu S, Pyatkevich Y. Ramsay Hunt syndrome type II. *Neurology* 2014;82:1664.
12. Bowman ME, Robinson JL. A child with ulcerative lesions and whitish plaques on the tongue. *Clin Infect Dis* 2010;51(1):77-8.
13. Neagu MR, Prasad S. Ramsay Hunt syndrome. *Pract Neurol* 2016;16(3):232.
14. Mangold AR, Torgerson RR, Rogers RS. Diseases of the tongue. *Clin Dermatol* 2016;34(4):458-69.
15. Braverman I, Uri N, Greenberg E. Trigeminal herpes zoster/chocolate-vanilla tongue. *Otolaryngol Head Neck Surg* 2000;122:463.
16. Shaikh S, Ta CN. Evaluation and management of herpes zoster ophthalmicus. *Am Fam Phys* 2002;66(9):1723-30.
17. Grose C, Bonthius D, Afifi AK. Chickenpox and the geniculate ganglion: facial nerve palsy, Ramsay Hunt syndrome, and acyclovir treatment. *Pediatr Infect Dis J* 2002;21:615-7.